



IBUPROFEN (Motrin, Advil)

Note: If possible, use weight to dose, otherwise use age. To arrive at the correct dose, weigh your child before giving Acetaminophen. A healthcare professional should be consulted for dosing in children under the age of two years.

	<i>Weight</i>	<i>6-11 lbs</i>	<i>12-17 lbs</i>	<i>18-23 lbs</i>	<i>24-35 lbs</i>	<i>36-47 lbs</i>	<i>48-59 lbs</i>	<i>60-71 lbs</i>	<i>72-95 lbs</i>
	<i>Age</i>	<i>0-5 mos</i>	<i>6-11 mos</i>	<i>12-23 mos</i>	<i>2-3 yrs</i>	<i>4-5 yrs</i>	<i>6-8 yrs</i>	<i>9-10 yrs</i>	<i>11 yrs</i>
<i>Concentrated Ibuprofen Infants' Drops</i> <i>Active Ingredient: Ibuprofen 50mg (in each 1.25 mL)</i>		---	<i>1.25 mL</i>	<i>1.875 mL</i>	---	---	---	---	---
<i>Children's Ibuprofen Suspension</i> <i>Active Ingredient: Ibuprofen 100mg (in each 5mL or 1 tsp)</i>		---	---	---	<i>1 tsp or 5mL</i>	<i>1 ½ tsp or 7.5mL</i>	<i>2 tsp or 10mL</i>	<i>2 ½ tsp or 12.5mL</i>	<i>3 tsp or 15mL</i>
<i>Ibuprofen Junior Strength Easy-to- Swallow Caplets</i> <i>Active Ingredient: Ibuprofen 100mg (in each caplet)</i>		---	---	---	---	---	<i>2 caplets</i>	<i>2 ½ caplets</i>	<i>3 caplets</i>

Take every 6-8 hours as needed. Do not exceed more than 4 doses in 24 hours.